


This article was downloaded by: [HEAL-Link Consortium]

On: 8 May 2009

Access details: Access Details: [subscription number 772810551]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Environmental Science and Health, Part C

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title-content=t713597270>

### 8-hydroxy-2' -deoxyguanosine (8-OHdG): A Critical Biomarker of Oxidative Stress and Carcinogenesis

ATHANASIOS VALAVANIDIS <sup>a</sup>; THOMAS VLACHOGIANNI <sup>a</sup>; CONSTANTINOS FIOTAKIS <sup>a</sup>

<sup>a</sup> Department of Chemistry, University of Athens, University Campus Zografou, Athens, Greece

Online Publication Date: 01 April 2009

**To cite this Article** VALAVANIDIS, ATHANASIOS, VLACHOGIANNI, THOMAS and FIOTAKIS, CONSTANTINOS(2009)'8-hydroxy-2' -deoxyguanosine (8-OHdG): A Critical Biomarker of Oxidative Stress and Carcinogenesis',Journal of Environmental Science and Health, Part C,27:2,120 — 139

**To link to this Article:** DOI: 10.1080/10590500902885684

**URL:** <http://dx.doi.org/10.1080/10590500902885684>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# 8-hydroxy-2'-deoxyguanosine (8-OHdG): A Critical Biomarker of Oxidative Stress and Carcinogenesis

Athanasios Valavanidis, Thomais Vlachogianni,  
and Constantinos Fiotakis

Department of Chemistry, University of Athens, University Campus Zografou, Athens, Greece

There is extensive experimental evidence that oxidative damage permanently occurs to lipids of cellular membranes, proteins, and DNA. In nuclear and mitochondrial DNA, 8-hydroxy-2'-deoxyguanosine (8-OHdG) or 8-oxo-7,8-dihydro-2'-deoxyguanosine (8-oxodG) is one of the predominant forms of free radical-induced oxidative lesions, and has therefore been widely used as a biomarker for oxidative stress and carcinogenesis. Studies showed that urinary 8-OHdG is a good biomarker for risk assessment of various cancers and degenerative diseases. The most widely used method of quantitative analysis is high-performance liquid chromatography (HPLC) with electrochemical detection (EC), gas chromatography-mass spectrometry (GC-MS), and HPLC tandem mass spectrometry. In order to resolve the methodological problems encountered in measuring quantitatively 8-OHdG, the European Standards Committee for Oxidative DNA Damage was set up in 1997 to resolve the artifactual oxidation problems during the procedures of isolation and purification of oxidative DNA products. The biomarker 8-OHdG or 8-oxodG has been a pivotal marker for measuring the effect of endogenous oxidative damage to DNA and as a factor of initiation and promotion of carcinogenesis. The biomarker has been used to estimate the DNA damage in humans after exposure to cancer-causing agents, such as tobacco smoke, asbestos fibers, heavy metals, and polycyclic aromatic hydrocarbons. In recent years, 8-OHdG has been used widely in many studies not only as a biomarker for the measurement of endogenous oxidative DNA damage but also as a risk factor for many diseases including cancer.

**Key Words:** Reactive oxygen species; biomarker; oxidative damage; DNA; 8-OHdG; 8-oxodG; carcinogenic substances; carcinogenesis

Received January 24, 2009; accepted March 11, 2009.

Address correspondence to Assoc. Prof. A. Valavanidis, Department of Chemistry, University of Athens University Campus Zografou, Athens, Greece. E-mail: valavanidis@chem.uoa.gr

## INTRODUCTION

Reactive oxygen species (ROS) are formed continuously in living cells of aerobic organisms as part of the physiological processes, metabolic, and other biochemical reactions. These endogenously produced ROS and oxygen-free radicals have important physiological functions, but because of their reactive nature can cause oxidative damage to lipids of cellular membranes, proteins, and DNA (1). Also, exogenous factors, such as UV radiation, tobacco smoke, asbestos, and carcinogenic substances, can produce ROS under various conditions (2).


Under normal physiological conditions in all aerobic organisms, there is a balance maintained between endogenous oxidants and numerous enzymatic and non-enzymatic antioxidant defenses (3,4). When an imbalance occurs, oxidants produce extensive oxidative damage to DNA, which, in turn, contributes to aging, malignant tumors, and other degenerative diseases (5–7).

In all living cells, damaged DNA is repaired enzymatically so that they regain their normal function, whereas misrepaired DNA can result in mutations (base substitution, deletions, and strand fragmentation) leading to carcinogenesis (8, 9). Although a broad range of DNA products are produced during oxidative damage to DNA (bases and sugar modifications, covalent crosslinks, single- and double-stranded breaks), most interest focused on nucleobase modifications and especially on the abundant lesion of 8-oxo-2'-deoxyguanosine because it is formed *in vivo* and can be measured quantitatively in cells following hydrolysis of the DNA to component bases (10).

### Reactive Oxygen Species and Oxidative DNA Damage

The most important oxygen-free radical causing damage to basic biomolecules (proteins, membrane lipids, and DNA) is the hydroxyl radical (HO•). The hydroxyl radical can be produced by various mechanisms, especially by the Fenton reaction of hydrogen peroxide (which diffuses into the nucleus) and metals and other endogenous and exogenous ROS. The HO• attacks DNA strands when it is produced adjacent to cellular and mitochondrial DNA causing the addition of DNA bases new radicals, which lead to the generation of a variety of oxidation products (11).

The interaction of HO• with the nucleobases of the DNA strand, such as guanine, leads to the formation of C8-hydroxyguanine (8-OHGua) or its nucleoside form deoxyguanosine (8-hydroxy-2'-deoxyguanosine). Initially, the reaction of the HO• addition leads to the generation of radical adducts, then by one electron abstraction, the 8-hydroxy-2'-deoxyguanosine (8-OH-dG) is formed (Fig. 1). The 8-OHdG undergoes keto-enol tautomerism, which favors the oxidized product 8-oxo-7,8-dihydro-2'-deoxyguanosine (8-oxodG). In the scientific literature 8-OHdG and 8-oxodG are used for the same compound (12).



**Figure 1:** Reaction of 2'-deoxyguanosine with hydroxyl radicals, radical adducts followed by reduction to 7-hydro-8-hydroxy-2'-deoxyguanosine, and by oxidation to 8-hydroxy-2'-deoxyguanosine (8-OHdG) or its tautomer 8-oxo-7-hydro-2'-deoxyguanosine (8-oxodG).

Although the other nucleobases of DNA react with  $\text{HO}\cdot$  in a similar manner, the 8-oxodG lesion is the most abundant DNA lesion because it is relatively easily formed and is promutagenic, and therefore a potential biomarker of carcinogenesis (12). Experiments showed that the mutagenic potential of 8-oxodG is supported by a loss of base-pairing specificity, misreading of adjacent

pyrimidines, or insertion of adenine opposite the lesions (13). Mutations that may arise due to the formation of 8-oxodG involving GC → TA transversion mutations (10, 14).

In recent years, the 8-OHdG lesions can be detected and analyzed with high sensitivity by high-performance liquid chromatography (HPLC), gas-chromatography-mass spectrometry (GC-MS), and liquid chromatography-mass spectrometry-mass spectrometry (LC-MS-MS) and by immunohistochemical methods and single cell gel electrophoresis. Determination and analysis of 8-OHdG can be performed in animal organs and in human samples (urine, human organs, leukocyte DNA) as a biomarker of oxidative stress, aging, and carcinogenesis (15).

### Key Stages of the 8-OHdG Discovery

The discovery of 8-OHdG was first reported in by Kasai and Nishimura in 1984 in their attempt to study and isolate mutagens in heated glucose (as a model of cooked food). Because of the difficulty to isolate mutagens, which are very unstable, they developed a method to trap reactive mutagens as guanine derivatives from the fact that carcinogens and mutagens react with nucleic acid bases, particularly guanine (16). The same authors found that oxygen-free radicals are involved in the C-8 oxidation reaction (17). In the following years, the formation of 8-OHdG was confirmed under exposures/reactions generating oxygen free radicals, such as asbestos fibers and H<sub>2</sub>O<sub>2</sub>. In the second case, H<sub>2</sub>O<sub>2</sub> and ferrous ions used a reducing agent (Fenton reaction) (18, 19) as did the carcinogenic cigarette smoke and diesel exhaust particles (20, 21)

In 1986, Floyd and coworkers (22) reported an analytical method for sensitive detection for 8-OHdG in cellular DNA by HPLC with electrochemical detector (HPLC-EC). The method was used by various scientific groups for numerous analytical studies of DNA damage in animal organs after the administration of various carcinogenic chemical substances and tumor promoters (23–26).

Similarly, numerous papers in recent years have analyzed levels of 8-OHdG in human organs and leukocyte DNA and in urine in relation to oxidative stress, diet, cancer incidence, and aging. As a result of these studies 8-OHdG has been established as an important biomarker of oxidative stress (12, 27), of cancer risk to humans by mechanisms of oxygen-free radicals (28, 29), of aging processes including degenerative diseases (30, 31), and in general as a biological marker of lifestyle and the effect of diet (32, 33).

In the past 20 years, numerous papers appeared in the scientific literature on the subject of oxidative stress and oxidative DNA damage, the role of oxygen-free radicals and ROS, and 8-OHdG or 8-oxodG. A search of Scholar Google in the end of 2008 showed 56,300 papers on oxidative DNA damage, 6,180 for 8-OHdG, and 2,430 for 8-oxodG.

## Methodology and Problems for the Analysis of 8-OHdG

The first reported analysis of 8-OHdG as a major and ubiquitous oxidation product of DNA in experimental animals in the urine and in humans was performed in 1989 (27). Since then several scientific groups have reported the 8-OHdG (or 8-oxodG) analysis by various methods.

The methodology of analysis followed two different strategies: (1) the direct approach designated to single out the DNA lesion by using physical and chemical methods and final DNA extraction and hydrolysis, and (2) the indirect approach, by preserving the whole DNA structure and the formation of the lesions are monitored *in situ* (34). In the first approach the quantitative analysis is performed by HPLC coupled with electrochemical detection (HPLC-EC), gas chromatography-mass spectrometry (GC-MS), and by HPLC tandem mass spectrometry (HPLC-MS/MS) (35–37). In the second approach the measurement is performed either by using antibodies that generally exhibit low specificity or through nicking activity of a specific DNA repair enzyme (38). Also, by using the enzyme formamidopyrimidine DNA N-glycosylase, the 8-oxodG is converted into a strand break and the quantitative measurement of the strand breaks is estimated by the comet assay, the alkaline elution technique, or the unwinding method (39–41).

Methodological differences among scientific laboratories directly analyzing the levels of 8-oxodG have resulted in a lack of consistency in the results, especially in the exaggeration of the background levels of 8-OHdG in human cellular DNA, sometimes by up to two or three orders of magnitude (42, 43). Although there are variations of levels between cell types and animal organs, such large differences were attributed to inaccurate experimental protocols and isolation methodologies that give rise to overestimations (probably by artefactual DNA oxidation during the procedure). As a general trend, levels of 8-oxodG obtained by the indirect approach (enzymatic assays) were found significantly lower than those of the direct methods of measurements (34).

The European Standards Committee on Oxidative Damage was set up in 1997 to resolve these issues (supported by the European Commission as a Concerted Action) with 27 analytical laboratories as members. The participating laboratories were provided with standard solutions of 8-oxoGua/8-oxodG in samples of calf thymus DNA, which required hydrolysis either to nucleosides using DNA-degrading enzymes (prior to analysis by HPLC) or to bases with formic acid (procedure normally used for GC-MS analysis) (44). Three samples were analyzed by all laboratories in triplicate by HPLC, GC-MS, and LC-MS-MS against their own standard. The results showed that the analysis of 8-oxodG in standard solutions was more accurate than previous analyses (15). The method LC-MS-MS was found to be demonstrably as reliable, sensitive, and precise as the best HPLC procedures and had the added advantage of giving unambiguous information on the identity of analytes. The HPLC

with electrochemical detection (EC-coulometric version) is more sensitive than the EC-amperometric version. The Committee noticed some problems with the GC-MS method, which resulted in higher values of 8-oxoGua as a result of oxidation occurring during sample preparation.

The direct approach of measurements, which requires isolation from cells of tissues, followed by hydrolysis and the release of the DNA lesions, has to be quantified by very sensitive detection techniques at the output of a chromatographic column. Thus, in each of the individual steps, artifactual DNA oxidation may occur, leading to overestimated levels (45, 46).

Cadet and colleagues (34) used a series of cell treatments methods, various digestion protocols, and analytical techniques to establish the cellular background level of 8-oxodG in cells. Their results showed that the steady-state cellular background level of 8-oxodG, in a lymphocyte cell line, was  $\sim 0.5$  lesions/ $10^6$  DNA nucleosides.

The levels of 8-oxodG were measured in DNA and compared under different methods, using HPLC-EC, EPLC-MS/MS, and HPLC-GC-MS analytical techniques (47–49). In recent years more sensitive analytical techniques used for the measurements and background levels of 8-oxodG in human urine were found at around 1 lesion/ $10^6$  DNA nucleosides (50–51)

## ENDOGENOUS OXIDATIVE DNA DAMAGE

### The Levels of 8-OHdG and Endogenous Oxidative DNA Damage

It was known that cellular (cDNA) and mitochondrial DNA (mtDNA) in aerobic organisms are constantly damaged even in the absence of any exposure to genotoxic carcinogens. Damaging processes include endogenous sources, such as DNA instability (e.g., depurination), spontaneous errors during DNA replication and repair, numerous physiological ROS as a result of oxygen metabolism, and products of lipid peroxidation. Exogenous sources also contribute to DNA damage, such as ionizing and UV radiation, naturally occurring radioisotopes, numerous genotoxic phytochemicals, and food pyrolysis products in diet and contaminants of air and water. Although aerobic biological systems are protected by antioxidant enzymatic and nonenzymatic antioxidants and by special DNA repair mechanisms that reduce substantially the levels of damage, a certain steady-state level is measurable as a background at all times (53).

Attempts to measure and quantify levels of background DNA damage were made in the 1980s (54, 55). The remarkable progress made in the next 10 years with respect to analytical sensitivity, structural characterization, and quantification of DNA lesions resolved many aspects of the problem. Oxidative stress, oxygen-free radicals, and lipid peroxidation products as byproducts of aerobic

metabolism were studied as genotoxic agents of endogenous DNA damage and were presented in a special issue of *Mutation Research* (56–59).

The etiology of various types of cancer is the result of DNA modifications and replication errors in somatic cells that give rise to mutations and subsequently to promotion and development to the malignant tumor. The basal steady-state levels of oxidative DNA base modifications (such as 8-oxodG) can be found in all types of cells, mostly from the continuous generation of ROS in the cellular oxygen metabolism (60). Scientists have long suspected that this endogenous basal DNA damage might play an important role in the initiation of carcinogenesis, whereas the environmental-exogenous effects are thought to mostly influence the stage of tumor promotion. The high incidence of sporadic cancer that is actually seen in our population can hardly be explained by the level of exposure to exogenous carcinogens of environmental pollution (61, 62). The hypothesis of the role of endogenous DNA damage by ROS attracted lots of attention by other scientists (8, 63, 64).

The correlation of the steady-state levels of oxidative DNA damage (ratio of the endogenous generation rate of DNA damage and the rate of the repair of these modifications, taking into account the cellular antioxidants and the efficiency of the repair system) with the mutation frequencies and the cancer incidence depended on the availability of sensitive methods of analysis of low levels of DNA damage. Since the 8-oxodG was the most frequently determined biomarker for the oxidative DNA damage its quantification at low levels was very important to test the model for the correlation (65).

The magnitude of endogenous DNA damage by ROS may be enormous. Each human cell metabolizes approximately  $10^{12}$  molecules of oxygen per day with 1% of oxygen metabolism, resulting in the generation of ROS (66). Based on the urinary excretion of characteristic modified nucleosides produced by ROS damage to DNA (8-oxodG and thymine glycol), it has been estimated that approximately 20,000 nucleobases in DNA are damaged in each human cell per day (67). The high level of oxidative damage by endogenous ROS was supported by Ames and colleagues, especially in the case of micronutrient deficiencies in humans (68, 69).

Although some scientists were doubtful as to the high levels of basal oxidative DNA damage, the quantitative estimates of steady-state levels of 8-oxodG, determined by various techniques in mammalian cells, ranged over several levels of magnitude. We now know that this was due to discrepancies in experimental procedures. The most recent reported estimates of steady-state levels of 8-oxodG in the DNA of human lymphocytes by various techniques were in the range of 0.2–2.0 lesions detected per  $10^6$  nucleobases (47).

### Relevance of Endogenous DNA Damage in Carcinogenesis

The relevance of endogenous DNA damage by ROS in carcinogenesis is a very difficult problem, and scientists tried to devise methods to estimate its



contribution. The *in vivo* studies have to overcome the additional influence of inflammation, which increases the level of oxidative damage and contributes to the increase of progression to cancer (64, 70). Also, epigenetic effects by ROS and oxidative stress, such as various types of signal transduction mechanisms, lead to apoptosis, modulate cell proliferation, and immune response, thus complicating the estimation of risk (71–73). Studies with antioxidants, such as vitamin C,  $\beta$ -carotene, and  $\alpha$ -tocopherol, for *in vivo* experiments showed inconsistent results. Dietary supplementation of the various antioxidants does not necessarily improve the antioxidant capacity of the cells and effect the steady-state levels of 8-oxoG or 8-oxodG in human lymphocytes in a human volunteer study (74) or in the liver of guinea-pigs (75).

Another approach of scientific research to solve the problem was to study the influence of DNA repair mechanisms on modulation of oxidative DNA damage. It is known that DNA base modifications are recognized and removed in eukaryotic cells by specific glycosylases that initiate the base excision repair pathway. For the lesion 8-oxoG, the most important and relevant enzyme in mammalian cells is the Ogg1 protein, in which the gene was cloned (76, 77). Overexpression of Ogg1 in cultured cells increased the repair rate by several fold but did not reduce the steady-state level or the spontaneous mutation rate significantly (78). Various scientific groups used the *ogg1*<sup>-/-</sup> knockout mice as a tool to analyze the effect of decreased DNA repair rates *in vivo*. Studies showed that there is an increase of the steady-state levels of 8-oxoG with the resulting increases of the spontaneous mutation frequencies observed in the liver of *ogg1*<sup>-/-</sup> mice (79–81). Results showed an increase of less than five additional 8-oxoG residues per 10<sup>6</sup> bp (double the spontaneous mutation rate). Therefore, 8-oxoG seems to play a major role for spontaneous mutagenesis (60).

The contribution of mitochondria to endogenous DNA damage of the nucleus, where the electron respiratory transport chain is the most important source of ROS in mammalian cells, was investigated by Epe and colleagues. The results showed that the contribution of mitochondria to the endogenously generated background levels of oxidative damage in the nuclear DNA is negligible (82).

## PRACTICAL APPLICATIONS OF THE 8-OHdG BIOMARKER

### Measurements of the 8-OHdG as a Biomarker in Relation to Exposures to Carcinogenic Substances

Analysis of urinary 8-OHdG has been established as an important biomarker to evaluate oxidative stress and to assess risk to cancer after exposure to various carcinogenic substances, environmental pollutants, and lifestyle factors (32, 83).

Tobacco smoking is well known for its oxidative properties and its carcinogenic potential (84). Elevated levels of 8-oxodG have been found in human tissues, including lungs and peripheral leukocytes of smokers (85, 86). Also, increased levels of 8-OHdG have been found in passive smokers in the workplace exposed to environmental tobacco smoke (87). Environmental tobacco smoke (ETS) can be a possible cancer risk factor in offspring of experimental animals. Exposure of pregnant rats to a relevant dose of 1 mg/m<sup>3</sup> of ETS during gestation resulted in a significant increase in 8-oxodG in maternal liver and in fetal liver and brain (88). The urinary excretion of products of damaged nucleotides in cellular pools or in DNA has been proved to be an important biomarker of risk for lung cancer. A cohort of 53,689 (men and women) aged 50–64 years, with 3–7 years follow-up, including 260 cases with lung cancer, were studied. Urinary excretion of 8-oxodG was higher in current smokers. Overall the incidence rate ratio (95% confidence interval) of lung cancer was 0.99 (0.80–1.22) per doubling of 8-oxodG excretion and there was no interaction with OGG1 genotype (89).

Asbestos fibers have been proved to be genotoxic in the lungs and are a potential carcinogenic hazard to occupationally exposed workers. Preliminary studies in cell free systems and in cultured cells showed that asbestos fibers produce hydroxyl radicals and cause oxidative DNA damage and act synergistically with cigarette smoke (90, 91). Other studies showed that inhaled asbestos fibers induce the formation of 8-OHdG in the DNA of white blood cells of workers highly exposed at the workplace (92, 93); high levels were found in the urine of asbestos-exposed workers, which correlated positively with duration of exposure (94).

Diesel exhaust particles and fine particulates, a mixture of organic chemicals, heavy metals, and carbon particles with persistent free radicals, are known by various studies to have the ability to cause oxidative injury and are potentially carcinogenic. Studies in mice showed a positive correlation of levels of 8-OHdG and lung carcinogenesis (95). In recent years many studies were devoted to studies of the importance of levels of urinary 8-OHdG in human lung tissues of workers exposed to diesel fumes or oil fly ash (96–98).

Ambient particulate air pollution in urban environments has been associated with an increase in lung cancer and cardiopulmonary mortality (99). The role of free radicals and other ROS formed by particulate matter (PM) has been linked to their increased toxicity in lung tissues (100). The biomarker 8-OHdG was used to study the effects of hydroxyl radical (HO•) generated both by coarse and fine PM (especially PM with aerodynamic diameter 10–2.5 μm, PM<sub>10</sub> and PM<sub>2.5</sub>) due to the heavy metals adsorbed in the pores and surfaces of the particles (101). Heavy metal content (Cu, Fe, V, etc.) of particulate matter and their bioreactivity was established as an important contributory factor to the enhancement of oxidative DNA damage (102, 103).

There are limited studies on the enhancement of urinary excretion of 8-oxodG or in the nasal respiratory epithelium in human subjects exposed to increased urban particulate air pollution, such as bus drivers and students working and living in central Copenhagen, children in Mexico City, and female toll station workers in Taipei (Taiwan) (104–107).

The correlation of inhaled particulate air pollution and oxidative DNA damage mechanisms by ROS (measured as 8-oxodG) or by stimulating cellular oxidant generation, has been recently reviewed (108–110).

### *Heavy metals*

Heavy metals and some metalloids are known to be carcinogenic to humans (111). The oxidative concept in metal carcinogenesis proposes the involvement of redox mechanisms, the production of hydroxyl radicals, which can attack DNA, causing hydroxylation and oxidative DNA damage. In turn, some of these products of DNA damage (including 8-oxodG) induce mutations leading to neoplastic transformations (112). Many studies investigated the carcinogenic potential of various metal ions via ROS generation (113, 114) and the induction of 8-OHdG lesion (115–117).

Polycyclic aromatic hydrocarbons (PAHs) have been known from epidemiological studies to be carcinogenic to humans. Elevated levels of 8-OHdG in urine or white blood cells of workers exposed to PAHs were determined for different working environments (e.g., coke-oven, roofers exposed to coal-tar pitch dust, etc) (118–121).

Various studies have investigated the effects of known carcinogens, such as benzene, styrene, and inorganic arsenic, using the urinary biomarker of 8-oxodG in workers exposed separately in different types working environments. The results of these studies showed a dose-response effect (122–124). A comprehensive review of the use of 8-OHdG as a marker of oxidative DNA damage in occupational and environmental exposures to carcinogenic substances was recently published (125).

## **Epidemiological Evidence for Carcinogenic Risk and 8-OHdG**

In recent years, the application of the biomarker 8-OHdG or 8-oxodG in molecular epidemiology studies as a risk factor for many diseases (especially cancer) associated with oxidative stress mechanisms is increasing (126). In the following section we present selected results only for recent years.

Elevated levels of urinary levels of 8-OHdG has been detected in patients with various malignancies. Chinese scientists measured urinary 8-OHdG in children with acute leukemia; their results suggested that elevated levels were associated with oxidative DNA damage. Furthermore, it was found that the risk was correlated with high levels of Cr exposure (127). Serum 8-OHdG was significantly increased in patients with colorectal cancer compared with

the control group. Also, the authors found that the activities of antioxidant enzymes in the study group were also significantly reduced (128). A recent study from Japan investigated the increased production of ROS and oxidative DNA damage in relation to hepatocarcinogenesis. Multivariate analysis found that levels of 8-OHdG and fibrosis were significant risk factor for hepatocellular carcinoma, especially in patients with hepatitis C virus infection (129).

Salivary analysis of 8-OHdG biomarker in patients with oral squamous cell carcinoma showed an increase of 65% compared with controls (130). Weight loss has been shown to increase oxidative DNA damage, thus increasing risk of some cancer forms, including lung cancer. An epidemiological study with 174 healthy employees showed that one unit decrease of Body Mass Index was associated with a 2.7% increase in 8-OHdG levels (131). Urinary 8-OHdG levels were also significantly higher among breast cancer patients than among control subjects, after making adjustments for confounders such as coffee consumption and use of oral contraceptives (132). Oxidative DNA damage, in the form of 8-OHdG levels, was found elevated for patients with Barrett's oesophagus and adenocarcinoma of oesophagus (133)

## CONCLUSIONS

Reactive oxygen species, including oxygen and nitrogen-free radicals, can cause specific oxidative DNA damage and play a leading role in initiation and promotion of carcinogenesis. The biomarker 8-OHdG or 8-oxodG, which was first reported by Kasai and Nishimura in 1984, has been established as a commonly measured and sensitive marker of DNA damage due to the hydroxyl radical attack at the C8 position of the nucleobase guanine or its nucleoside guanosine. Results showed that this damage, if left unrepaired, may contribute to mutagenicity and cancer promotion. Many studies in the past 20 years and improvements in the quantitative estimation of 8-OHdG by various analytical techniques in blood cells or in urine have established it as a very important biomarker not only for carcinogenesis but also for aging and degenerative diseases.

## REFERENCES

1. Kehrer JP. Free radicals as mediators of tissue injury and disease. *Critical Reviews in Toxicology* 1993;23:21–48.
2. Dröge W. Free radicals in the physiological control of cell function. *Physiology Reviews* 2002;82:47–95.
3. Scandalios JG (Ed.). *Oxidative Stress and the Molecular Biology of Antioxidant Defenses*. New York: Cold Spring Harbor Laboratory Press; 1997.
4. Halliwell B, Gutteridge JMC. *Free Radical in Biology and Medicine*, 3rd ed. Oxford: Oxford University Press; 1999.

5. Hartman D. Free Radicals, Ageing and Degenerative Disease. New York: Alan Liss Inc; 1986.
6. Poulsen HE, Prieme S, Loft S. Role of oxidative DNA damage in cancer initiation and promotion. *European Journal of Cancer Prevention* 1998;7:9–16.
7. Beckman KB, Ames BN. The free radical theory of aging matures. *Physiology Reviews* 1998;78:547–581.
8. Loft S, Poulsen HE. Cancer risk and oxidative DNA damage in man. *Journal of Molecular Medicine* 1996;74:297–312.
9. Kawanishi S, Hiraku Y, Oikawa S. Mechanism of guanine-specific DNA damage by oxidative stress and its role in carcinogenesis and aging. *Mutation Research* 2001;488:65–76.
10. Lunec J, Holloway KA, Cooke MS, Faux S, Griffiths HR, Evans MD. Urinary 8-oxo-2'-deoxyguanosine: Redox regulation of DNA repair in vivo? *Free Radical Biology and Medicine* 2002;33:875–885.
11. Valko M, Izakovic M, Mazur M, Rhodes CJ, Telser J. Role of oxygen radicals in DNA damage and cancer incidence. *Molecular and Cellular Biochemistry* 2004;266:37–56.
12. Kasai H. Analysis of a form of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis. *Mutation Research* 1997;387:147–163.
13. Kasai H. Chemistry-based studies on oxidative DNA damage: Formation, repair, and mutagenesis. *Free Radical Biology and Medicine* 2002;33:450–456.
14. Cheng KC, Cahill DS, Kasai H, Nishimura S, Loeb LA. 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G→T and A→C. substitutions. *Journal of Molecular Biology* 1992;267:166–172.
15. European Standard Committee on Oxidative DNA Damage. Comparison of different methods of measuring 8-oxoguanine as a marker of oxidative DNA damage. *Free Radical Biology and Medicine* 2000;32:333–341.
16. Kasai H, Hayami H, Yamaizumi Z, Saito H, Nishimura S. Detection and identification of mutagens and carcinogens as their adducts with guanosine derivatives. *Nucleic Acids Research* 1984;12:21270–2136.
17. Kasai H, Nishimura S. Hydroxylation of deoxyguanosine at the C-8 position by polyphenols and aminophenols in the presence of hydrogen peroxide and ferric ion. *Gann (Japanese Journal of Cancer Research)* 1984;75:565–566.
18. Kasai H, Nishimura S. DNA damage induced by asbestos in the presence of hydrogen peroxide. *Gann* 1984;75:841–844.
19. Kasai H, Nishimura S. Hydroxylation of deoxyguanosine at the C-68 position by ascorbic acid and other reducing agents. *Nucleic Acids Research* 1984;12:2137–2145.
20. Leaderson P, Tagesson C. Cigarette smoke-induced DNA-damage: role of hydroquinone and catechol in the formation of the oxidative DNA-adduct, 8-hydroxydeoxyguanosine. *Chemical–Biological Interactions* 1990;75:71–81.
21. Nagashima M, Kasai H, Yokota J, Nagamachi T, Ichinose T, Sagai M. Formation of an oxidative DNA damage, 8-hydroxydeoxyguanosine, in mouse lung DNA after intracheal instillation of diesel exhaust particles and effects of high dietary fat and beta-carotene on this process. *Carcinogenesis* 1995;16:1441–1445.

22. Floyd RA, Watson JJ, Wong PK, Altmiller DH, Rickard RC. Hydroxyl free radical adduct of deoxyguanosine: sensitive detection and mechanisms of formation. *Free Radical Research and Communications* 1986;1:163–172.
23. Fiala ES, Conaway C, Mathis JF. Oxidative DNA and RNA damage in the livers of Sprague-Dawley rats treated with the hepatocarcinogen 2-nitropropane. *Cancer Research* 1989;49:5518–5522.
24. Umemura T, Sai K, Takagi A, Hasegawa R, Kurokawa Y. Oxidative DNA damage, lipid peroxidation and nephrotoxicity induced in the rat kidney after ferric nitrilotriacetate administration. *Cancer Letters* 1990;54:95–100.
25. Tagaki T, Sai K, Umemura T, Hasegawa R, Kurokawa Y. Short-term exposure to peroxisome proliferators, perfluorooctanoic acid and perfluorodecanoic acid, causes significant increase of 8-hydroxydeoxyguanosine in liver DNA of rats. *Cancer Letters* 1991;57:55–60.
26. Cattley RC, Glover SE. Elevated 8-hydroxydeoxyguanosine in hepatic DNA of rats following exposure to peroxisome proliferators: relationship to carcinogenesis and nuclear localization. *Carcinogenesis* 1993;14:2495–2499.
27. Shigenaga MK, Gimeno CJ, Ames BN. Urinary 8-hydroxy-2'-deoxyguanosine as a biological marker of *in vivo* oxidative DNA damage. *Proceedings National Academy of Sciences, USA* 1989;86:9697–9701.
28. Halliwell B. Can oxidative DNA damage be used as a biomarker of cancer risk in humans? Problems, resolutions, and preliminary results from nutritional supplementation studies. *Free Radical Research* 1998;29:469–486.
29. Marnett LJ. Oxyradicals and DNA damage. *Carcinogenesis* 2000;21:361–370.
30. Fraga CG, Shigenaga MK, Park JW, Degan P, Ames BN. Oxidative damage to DNA during aging: 8-hydroxy-2'-deoxyguanosine in rat organ DNA and urine. *Proceedings of National Academy of Sciences, USA* 1990;87:4533–4537.
31. Sayre LM, Smith MA, Perry G. Chemistry and biochemistry of oxidative stress in neurodegenerative disease. *Current Medicinal Chemistry* 2001;8:721–738.
32. Kasai H, Iwamoto-Tanaka N, Miyamoto T, Kawanami K, Kawanami S, Kido R, Ikeda M. Life style and urinary 8-hydroxydeoxyguanosine, a marker of oxidative DNA damage: effects of exercise, working conditions, meat intake, body mass index, and smoking. *Japanese Journal of Cancer Research* 2001;92:9–15.
33. Halliwell B. Effect of diet on cancer development: is oxidative DNA damage a biomarker? *Free Radical Biology and Medicine* 2002;32:986–974.
34. Ravanat J-L, Douki T, Duez P, Gremaud E, Herbert K, Hofer T, Lasserre L, Saint-Pierre C, Favier A, Cadet J. Cellular background level of 8-oxo-7,8-dihydro-2'-deoxyguanosine: an isotope based method to evaluate artefactual oxidation of DNA during its extraction and subsequent work-up. *Carcinogenesis* 2002;23:1911–1918.
35. Serrano J, Palmeira CM, Wallace KB, Kuehl DW. Determination of 8-hydroxyguanosine in biological tissues by liquid chromatography/electrospray ionization-mass spectrometry/mass spectrometry. *Rapid Communication Mass Spectrometry* 1996;10:1789–1791.
36. Dizdaroglu M, Jaruga P, Birincioglu M, Rodriguez H. Free radical-induced damage to DNA: mechanisms and measurement. *Free Radical Biology and Medicine* 2002;32:1102–1115.
37. Kasai H. A new automated method to analyze urinary 8-hydroxydeoxy-guanosine by high-performance liquid chromatography-electrochemical detector system. *Journal of Radiation Research* 2003;44:185–189.

38. Yin B, Whyatt RM, Perera FP, Randall MC, Cooper TB, Santella RM. Determination of 8-hydroxydeoxyguanosine by an immunoaffinity chromatography-monoclonal antibody-based ELISA. *Free Radical Biology and Medicine* 1995;18:1023–1032.
39. Collins AR, Duthie SJ, Dobson VL. Direct enzymic detection of endogenous oxidative base damage in human lymphocyte DNA. *Carcinogenesis* 1993;14:1733–1735.
40. Epe B, Hegler J. Oxidative DNA damage: endonuclease fingerprinting. *Methods in Enzymology* 1994;234:122–131.
41. Hartwig A, Dally H, Schlepegrell R. Sensitive analysis of oxidative DNA damage in mammalian cells: use of the bacterial Fpg protein in combination with alkaline unwinding. *Toxicology Letters* 1996;88:85–90.
42. Collins AR, Cadet J, Epe B, Gedik C. Problems in the measurement of 8-oxoguanine in human DNA. Report of a workshop, held in Aberdeen, UK, 19–21 January, 1997. *Carcinogenesis* 1997;18:1833–1836.
43. Cadet J, D'Ham C, Douki T, Pouget J-P, Ravanat J-L, Sauvaigo S. Facts and artifacts in the measurement of oxidative base damage to DNA. *Free Radical Research* 1998;29:541–550.
44. European Standard Committee on Oxidative DNA Damage. Inter-laboratory validation of procedures for measuring 8-oxo-7,8-dihydroguanine/8-oxo-7,8-dihydro-2'-deoxyguanosine in DNA. *Free Radical Research* 2002;36:239–245.
45. Ravanat J-L, Turesky RJ, Gremaud E, Trudel LJ, Stadler RH. Determination of 8-oxoguanine in DNA by gas chromatography-mass spectrometry and HPLC-electrochemical detection. Overestimation of the background level of the oxidized base by gas chromatography-mass spectrometry assay. *Chemical Research in Toxicology* 1995;8:1039–1045.
46. Finnegan MTV, Herbert KE, Evans MD, Griffiths HR, Lunec, J. Evidence for sensitization of DNA to oxidative damage during isolation. *Free Radical Biology and Medicine* 1996;20:93–98.
47. Frelon S, Douki T, Ravanat J-L, Pouget JP, Tornabene C, Cadet J. High performance liquid chromatography-tandem mass spectrometry for the measurement of radiation-induced base damage to isolated and cellular DNA. *Chemical Research in Toxicology* 2000;13:1002–1010.
48. Rodriguez H, Jurado J, Laval J, Dizdaroglu M. Comparison of the levels of 8-hydroxyguanine in DNA as measured by gas chromatography-mass spectrometry following hydrolysis of DNA by *Escherichia coli* Fpg protein or formic acid. *Nucleic Acids Research* 2000;29:E75.
49. Dizdaroglu M, Jaruga P, Rodriguez H. Measurements of 8-hydroxy-2'-deoxyguanosine in DNA by high-performance liquid chromatography-mass spectrometry: comparison with measurement by gas chromatography-mass spectrometry. *Nucleic Acids Research* 2001;29:E12.
50. Weinmann A, Belling D, Poulsen HE. Measurement of 8-oxo-2'-deoxyguanosine and 8-oxo-2'-deoxyadenosine in DNA and human urine by high-performance liquid chromatography-electrospray tandem mass spectrometry. *Free Radical Biology and Medicine* 2001;30:757–764.
51. Sabatini L, Barbieri A, Tosi M, Roda A, Violante FS. A method for routine quantification of urinary 8-hydroxy-2'-deoxyguanosine based on solid-phase extraction and micro-high-performance liquid chromatography/electrospray ionization tandem mass spectrometry. *Rapid Communication Mass Spectrometry* 2005;19:147–152.

52. Harri M, Kasai H, Mori T, Tornaues J, Savela K, Peltonen, K. Analysis of 8-hydro-2'-deoxyguanosine in urine using high-performance liquid chromatography-electrospray tandem mass spectrometry. *Journal of Chromatography B, Analytical. Technology. Biomedicine. Life Sciences* 2007;853:242–246.
53. Gupta RC, Lutz WK. Background DNA damage from endogenous and unavoidable exogenous carcinogens: a basis for spontaneous cancer incidence? *Mutation Research* 1999;424:1–8.
54. Saul RL, Ames BN. Background levels of DNA damage in the population. In: Simic MG, Grossman L, Upton AC (Eds.), *Mechanisms of DNA Damage*. New York: Repair Plenum Press; 1986: 529–535.
55. Lutz WK. Endogenous genotoxic agents and processes as a basis of spontaneous carcinogenesis. *Mutation Research* 1990;239:287–295.
56. Cadet J, Delatour T, Douki T, Gasparutto D, Pouget J-P, Ravanat J-L, Sauvaigo S. Hydroxyl radicals and DNA base damage. *Mutation Research* 1999;424:9–21.
57. Lloyd DR, Phillips DH. Oxidative DNA damage mediated by copper(II), iron(II) and nickel(II) Fenton reaction: evidence for site-specific mechanisms in the formation of double-strand breaks, 8-hydroxydeoxy-guanosine and putative intrastrand crosslinks. *Mutation Research* 1999;424:23–36.
58. Beckman KB, Ames BN. Endogenous oxidative damage of mtDNA. *Mutation Research* 1999;424:51–58.
59. Marnett LJ. Lipid peroxidation-DNA damage by malondialdehyde. *Mutation Research* 1999;424:83–95.
60. Epe B. Role of endogenous oxidative DNA damage in carcinogenesis: what can we learn from repair-deficient mice? *Biological Chemistry* 2002;383:467–475.
61. Gold LS, Slone TH, Stern BR, Manley NB, Ames BN. Rodent carcinogens: setting priorities. *Science* 1992;258:261–265.
62. Ames BN, Gold LS. Paracelsus to parascience: the environmental cancer distraction. *Mutation Research* 2000;447:3–13.
63. Collins AR. Oxidative DNA damage, antioxidants, and cancer. *Bioassays* 1999;21:238–246.
64. Jackson AL, Loeb LA. The contribution of endogenous sources of DNA damage to the multiple mutations in cancer. *Mutation Research* 2001;477:7–21.
65. Otteneeder M, Lutz WK. Correlation of DNA adduct levels with tumor incidence: carcinogenic potency of DNA adducts. *Mutation Research* 1999;424:237–247.
66. Floyd RA. Measurement of oxidative stress in vivo. In Davies, KJA, Ursini F (Eds.), *The Oxygen Paradox*. Cleup: Cleup University Press; 1995:89–103.
67. Beckman KB, Ames BN. Oxidative decay of DNA. *Journal of Biological Chemistry* 1997;272:19633–19636.
68. Richter C, Park J-W, Ames BN. Normal oxidative damage to mitochondrial and nuclear DNA is extensive. *Proceedings of National Academy of Sciences, USA* 1988;85:6465–6467.
69. Ames BN. DNA damage from micronutrient deficiencies is likely to be a major cause of cancer. *Mutation Research* 2001;475:7–20.
70. Wiseman H, Halliwell B. Damage to DNA by reactive oxygen and nitrogen species: role of inflammatory disease and progression to cancer. *Biochemical Journal* 1996;313:17–29.



71. Poulsen HE, Jensen BR, Weimann A, Jensen SA, Sorensen M, Loft S. Antioxidants, DNA damage and gene expression. *Free Radical Research* 2000;33:33–39.
72. Allen RG, Tresini M. Oxidative stress and gene regulation. *Free Radical Biology and Medicine* 2000;28:463–499.
73. Schoonbroodt S, Piette J. Oxidative stress interference with the nuclear factor  $\kappa$ B activation pathways. *Biochemical Pharmacology* 2000;60:1075–1063.
74. Anderson D, Phillips BJ, Yu TW, Edwards AJ, Ayes R, Butterworth KR. The effects of vitamin C supplementation on biomarkers of oxygen radical generated damage in human volunteers with “low” or “high” cholesterol levels. *Environmental Molecular Mutagenesis* 1997;30:161–174.
75. Cadenas S, Barja G, Poulsen HE, Loft S. Oxidative DNA damage estimated by 8-oxodG in the liver of guinea-pigs supplemented with graded dietary doses of ascorbic acid and  $\alpha$ -tocopherol. *Carcinogenesis* 1997;18:2373–2377.
76. Boiteux S, Radicella, JP. The human OGG1 gene: structure, functions, and its implication in the process of carcinogenesis. *Archives of Biochemistry and Biophysics* 2000;377:1–8.
77. Bruner SD, Norman PDG, Verdine YL. Structural basis for recognition and repair of the endogenous mutagen 8-oxoguanine in DNA. *Nature* 2000;403:859–866.
78. Hollenbach S, Dhenaut A, Eckert I, Radicella JP, Epe B. Overexpression of Ogg1 in mammalian cells: effects on induced and spontaneous oxidative DNA damage and mutagenesis. *Carcinogenesis* 1999;20:1863–1868.
79. Klungland A, Rosewell I, Hollenbach S, Larsen E, Daly G, Epe B, Seeberg E, Lindahl T, Barnes DE. Accumulation of premutagenic DNA lesions in mice defective in removal of oxidative base damage. *Proceedings of National Academy of Sciences, USA* 1999;96:13300–13305.
80. Minowa O, Arai T, Hirano M, Monden Y, Nakai S, Fukuda M, Itoh M. *Mmh/Ogg1* gene inactivation results in accumulation of 8-hydroxyguanine in mice. *Proceedings National Academy of Sciences, USA* 2000;97:4156–4161.
81. Osterod M, Hollenbach S, Hengstler JG, Barnes DE, Lindahl T, Epe B. Age-related and tissue-specific accumulation of oxidative DNA base damage in 7,8-dihydro-8-oxoguanine-DNA glycosylase (Ogg1) deficient mice. *Carcinogenesis* 2001;22:1459–1463.
82. Hoffmann S, Spitkovsky D, Radicella JP, Epe B, Wiesner, RJ. Reactive oxygen species derived from the mitochondrial respiratory chain are not responsible for the basal levels of oxidative base modifications observed in nuclear DNA of mammalian cells. *Free Radical Biology and Medicine* 2004;36:765–773.
83. Cooke MS, Lunec J, Evans MD. Progress in the analysis of urinary oxidative DNA damage. *Free Radical Biology and Medicine* 2002;33:1601–1614.
84. Pryor WA. Cigarette smoke radicals and the role of free radicals in chemical carcinogenesis. *Environmental Health Perspectives* 1997;105:875–882.
85. Kiyosawa H, Suko M, Okudaira H, Murata K, Miyamoto T, Chung M-H, Kasai H. Cigarette smoking induce formation of 8-hydroxydeoxy-guanosine, one of the oxidative DNA damages in human peripheral leukocytes. *Free Radic. Res. Commun.* 1990;11:23–27.
86. Asami S, Hirano T, Yamaguchi, R, Tomioka Y, Itoh H, Kasai H. Increase of a type of oxidative DNA damage, 8-hydroxyguanine, and its repair activity in human leukocytes by cigarette smoking. *Cancer Research* 1996;56:2546–2549.

87. Howard DJ, Ota RB, Briggs LA, Hampton M, Pritsos CA. Environmental tobacco smoke in the workplace induces oxidative stress in employees, including increased production of 8-hydroxy-2'-deoxyguanosine. *Cancer Epidemiology Biomarkers and Prevention* 1998;7:141–146.
88. Maciag A, Bialkowska A, Espiritu I, Powell D, Alvord WG, Kasprzak KS, Anderson LM, Witschi HR. Gestation stage-specific oxidative deoxyribonucleic acid damage from sidestream smoke in pregnant rats and their fetuses. *Archives of Environmental Health* 2003;58:238–244.
89. Loft S, Svoboda P, Kasai H, Tjønneland A, Vogel U, Møller P, Overvad K, Raaschou-Nielsen O. Prospective study of 8-oxo-7,8-dihydro-2'-deoxyguanosine excretion and the risk of lung cancer. *Carcinogenesis* 2006;27:1245–1250.
90. Jackson JH, Schraufstatter IU, Hyslop PA, Vosbeck K, Sauerheber R, Weitzman SA, Cochrane CG. Role of oxidants in DNA damage. Hydroxyl radical mediates the synergistic DNA damaging effects of asbestos and cigarette smoke. *Journal of Clinical Investigation* 1987;80:1090–1095.
91. Chen Q, Marsh J, Ames B, Mossman B. Detection of 8-oxo-2'-deoxyguanosine, a marker of oxidative DNA damage, in culture medium from human mesothelial cells exposed in crocidolite asbestos. *Carcinogenesis* 1996;17:2525–2527.
92. Marczynski B, Rozynek P, Kraus T, Schlosser S, Raithel HJ, Baur X. Levels of 8-hydroxy-2'-deoxyguanosine in DNA of white blood cells from workers highly exposed to asbestos in Germany. *Mutation Research* 2000;468:195–202.
93. Unfried K, Schurkes C, Abel J. Distinct spectrum of mutations induced by crocidolite asbestos: clue for 8-hydroxydeoxyguanosine dependent mutagenesis *in vivo*. *Cancer Research* 2002;62:99–104.
94. Yoshida R, Ogawa Y, Shioji I, Yu X, Shibata E, Mori I, Kubota H, Kishida A, Hisanaga N. Urinary 8-oxo-7,8-dihydro-2'-deoxyguanosine and biopyrrins levels among construction workers with asbestos exposure history. *Industrial Health* 2001;39:186–188.
95. Ichinose T, Yajima Y, Nagashima M, Takenoshita S, Nagamachi Y, Sagai M. Lung carcinogenesis and formation of 8-hydroxydeoxyguanosine in mice by diesel exhaust particles. *Carcinogenesis* 1997;18:185–192.
96. Tokiwa H, Sera N, Nakanishi Y, Sagai M. 8-Hydroxyguanosine formed in human lung tissues and the association with diesel exhaust particles. *Free Radical Biology and Medicine* 1999;27:1251–1258.
97. Kim JY, Mukherjee S, Ngo L, Christiani DC. Urinary 8-hydroxy-2'-deoxyguanosine as a biomarker of oxidative DNA damage in workers exposed to fine particulates. *Environmental Health Perspectives* 2004;112:666–671.
98. Harri M, Svoboda P, Mori T, Mutanen P, Kasai H, Savela K. Analysis of 8-hydroxydeoxyguanosine among workers exposed to diesel particulate exhaust: comparison with urinary metabolites and PAH air monitoring. *Free Radical Research* 2005;39:963–972.
99. Pope CA, III, Burnett RT, Thun MJ, Calle EE, Krewski D, Ito K, Thurston GD. Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *Journal of American Medical Association* 2002;287:1132–1141.
100. Dellinger B, Pryor WA, Cueto R, Squadrito GL, Hedge V, Deutsch W A. Role of free radicals in the toxicity of airborne fine particulate matter. *Chemical Research in Toxicology* 2001;14:1371–1377.

101. Shi T, Knaapen AM, Begerow J, Birmili W, Borm PJA, Schins RPF. Temporal variation of hydroxyl radical generation and 8-hydroxy-2'-deoxyguanosine formation by coarse and fine particulate matter. *Occupational and Environmental Medicine* 2003;60:315–321.
102. Prahalat AK, Inmon J, Ghio AJ, Gallagher JE. Enhancement of 2'-deoxyguanosine hydroxylation and DNA damage by coal and oil fly ash in relation to particulate metal content and availability. *Chemical research in Toxicology* 2000;13:1011–1019.
103. Prahalat AK, Inmon J, Dailey LA, Madden MC, Ghio AJ, Gallagher JE. Air pollution particles mediated oxidative DNA base damage in a cell free system and in human airway epithelial cells in relation to particulate metal content and bioreactivity. *Chemical Research in Toxicology* 2001;14:879–887.
104. Loft S, Poulsen HE, Vistisen LE. Increased urinary excretion of 8-oxo-2'-deoxyguanosine, a biomarker of oxidative DNA damage, in urban bus drivers. *Mutation Research* 1999;441:11–19.
105. Sørensen M, Autrup H, Hertel O, Wallin H, Knudsen LE, Loft S. Personal exposure to PM2.5 and biomarkers of DNA damage. *Cancer Epidemiology Biomarkers Prevention* 2003;12:191–196.
106. Calderon-Carciduenas L, Wen-Wang L, Zhang YJ, Rodriguez-Alcaraz A, Osnaya N, Villarreal-Calderon A, Santella RM. 8-hydroxy-2'-deoxyguanosine, a major mutagenic oxidative DNA lesion and DNA strand breaks in nasal respiratory epithelium of children exposed to urban pollution. *Environmental Health Perspectives* 1999;107:469–474.
107. Lai CH, Liou SH, Lin HC, Shih TS, Tsai FJ, Chen JS, Yang T, Jaakkola JJ, Strickland PT. Exposure to traffic exhausts and oxidative DNA damage. *Occupational and Environmental Medicine* 2005;62:216–222.
108. Knaapen AM, Borm PJA, Albrecht C, Shins RPF. Inhaled particles and lung cancer. Part A: mechanisms. *International Journal of Cancer* 2004;109:799–809.
109. Risom, L. Moller P, Loft S. Oxidative stress-induced DNA damage by particulate air pollution. *Mutation Research* 2005;592:119–137.
110. De Kok TM, Drieste HA, Hogervorst JG, Briedé JJ. Toxicological assessment of ambient and traffic-related particulate matter: a review of recent studies. *Mutation Research* 2006;613:103–122.
111. Kasprzak KS. Oxidative DNA and protein damage in metal-induced toxicity and carcinogenicity. *Free Radical Biology and Medicine* 2002;32:958–967.
112. Bal W, Kasprzak KS. Induction of oxidative DNA damage by carcinogenic metals. *Toxicology Letters* 2002;127:55–62.
113. Toyokumi S, Sagripanti JL. Association between 8-hydroxy-2'-deoxyguanosine formation and DNA strand breaks mediated by copper and iron. *Free Radical Biology and Medicine* 1996;20:859–864.
114. Shi X, Jiang H, Mao Y, Ye J, Saffiotti U. Vanadium (IV)-mediated free radical generation and related 2'-deoxyguanosine hydroxylation and DNA damage. *Toxicology* 1996;106:27–38.
115. Mao Y, Liu KJ, Jiang JJ, Shi X. Generation of reactive oxygen species by Co(II) from H2O2 in the presence of chelators in relation to DNA damage and 2'-deoxyguanosine hydroxylation. *Journal of Toxicology and Environmental Health* 1996;47:61–75.

116. Kuo HW, Chang SF, Wu KY, Wu FY. Chromium(VI) induced oxidative damage to DNA: increase of urinary 8-hydroxydeoxyguanosine concentrations (8-OHdG) among electroplating workers. *Occupational and Environmental Medicine* 2003;60:590–594.
117. Valavanidis A, Vlahoyianni T, Fiotakis K. Comparative study of the formation of oxidative DNA damage marker 8-hydroxy-2'-deoxyguanosine (8-OHdG) adduct from the nucleoside 2'-deoxyguanosine by transition metals and suspensions of particulate matter in relation to metal content and redox reactivity. *Free Radical Research* 2005;39:1071–1081.
118. Toraason M, Hayden C, Marow D, Rinehart R, Mathgias P, Werren D, DeBord DG, Reid TM. DNA strand breaks, oxidative damage, and 1-OH pyrene in roofers with coal-tar pitch dust and/or asphalt fume exposure. *International Archives Occupational and Environmental Health* 2001;74:396–404.
119. Marczyński B, Rihs HP, Rossbach B, Hölzer J, Angerer J, Scherenberg M, Hoffmann G, Brünnig T, Wilhelm M. Analysis of 8-oxo-7,8-dihydro-2'-deoxyguanosine and DNA strand breaks in white blood cells of occupationally exposed workers: comparison with ambient monitoring, urinary metabolites and enzyme polymorphisms. *Carcinogenesis* 2002;23:273–281.
120. Zhang J, Ichiba M, Hanaoka T, Pan G, Yamano Y, Hara K, Takahashi K, Tomokuni K. Leukocyte 8-hydroxydeoxyguanosine and aromatic DNA adduct in coke-oven workers with polycyclic aromatic hydrocarbon exposure. *International Archives of Occupational and Environmental Health* 2003;76:499–504.
121. Marczyński B, Preuss R, Mensing T, Angerer J, Seidel A, El Mourabit A, Wilhelm M, Brüning T. Genotoxic risk assessment in white blood cells of occupationally exposed workers before and after alteration of the polycyclic aromatic hydrocarbon (PAH) profile in the production material: comparison with PAH air and urinary metabolite levels. *International Archives of Occupational and Environmental Health* 2005;78:97–108.
122. Lagorio S, Tagesson C, Forastiere F, Iavarone I, Axelson O, Carere A. Exposure to benzene and urinary concentrations of 8-hydroxydeoxy-guanosine, a biological marker of oxidative damage to DNA. *Occupational and Environmental Medicine* 1994;51:739–743.
123. Marczyński B, Rozynek P, Elliehausen HJ, Korn M, Baur X. Detection of 8-hydroxydeoxyguanosine, a marker of oxidative DNA damage, in white blood cells of workers occupationally exposed to styrene. *Archives of Toxicology* 1997;71:496–500.
124. Hu C-W, Pan C-H, Huang Y-H, Wu M-T, Chang LW, Wang C-J, an Chao M-R. Effects of arsenic exposure among semiconductor workers: a cautionary note on urinary 8-oxo-7,8-dihydro-2'-deoxyguanosine. *Free Radical Biology and Medicine* 2006;40:1273–1278.
125. Pilger A, Rüdiger HW. 8-Hydroxy-2'-deoxyguanosine as a marker of oxidative DNA damage related to occupational and environmental exposures. *International Archives of Occupational and Environmental Health* 2006;80:1–15.
126. Toyokuni S. Molecular mechanisms of oxidative stress-induced carcinogenesis: from epidemiology to oxygenomics. *International Union of Biochemistry and Molecular Biology, Life* 2008;60:441–447.
127. Yang Y, Tian Y, Yan C, Jin X, Tang J, Shen X. Determinants of urinary 8-hydroxy-2'-deoxyguanosine in Chinese children with acute leukemia. *Environmental Toxicology* 2008 [Oct 31 Epub ahead of print].
128. Chang D, Wang F, Zhao YS, Pan HZ. Evaluation of oxidative stress in colorectal cancer patients. *Biomedical Environmental Science* 2008;21:286–289.

129. Chuma M, Hige S, Nakanishi M, Ogawa K, Natsuizaka M, Yamamoto Y, Asaka M. 8-Hydroxy-2'-deoxy-guanosine is a risk factor for development of hepatocellular carcinoma in patients with chronic hepatitis C virus infection. *Journal of Gastroenterology* 2008;23:1431–1436.
130. Bahar G, Felnmesser R, Shpitzer T, Popovtzer A, Nagler RM. Salivary analysis in oral cancer patients: DNA and protein oxidation, reactive nitrogen species, and antioxidant profile. *Cancer* 2007;109:54–59.
131. Mizoue T, Tokunaga S, Kasai H, Kawai K, Sato M, Kubo T. Body mass index and oxidative DNA damage: a longitudinal study. *Cancer Science* 2007;98:1254–1258.
132. Kuo HW, Chou SY, Hu TW, Wu FY, Chen DJ. Urinary 8-hydroxy-2'-deoxyguanosine (8-OHdG) and genetic polymorphisms in breast cancer patients. *Mutation Research* 2007;631:62–68.
133. Räsänen JV, Sihvo EI, Ahotupa MO, Farkkila MA, Salo JA. The expression of 8-hydroxydeoxyguanosine in oesophageal tissues and tumours. *European Journal of Surgical Oncology* 2007;33:1164–1168.